

SIMBOTIN, G., ing.; TUDOROIU, V.; MIHAILESCU, E., ing.; BRANA, G.

Way of organizing new systems of industrial telemechanics.  
Automatica electronica 7 no.3:106-110 My-Je '63.

1. Cercetator principal la Institutul de Cercetari Electrotehnice  
(for Simbotin, Tudoroiu). 2. Cercetator stagiar la Institutul  
de Cercetari Electrotehnice (for Mihailescu, Brana).

"APPROVED FOR RELEASE: 03/14/2001

CIA-RDP86-00513R001757410002-4

APPROVED FOR RELEASE: 03/14/2001

CIA-RDP86-00513R001757410002-4"

TUDOROIU, V.

Automation in the extraction, transportation, and distribution of methane gas. p. 104.

AUTOMATICA SI ELECTRONICA (Asociatia Stinifica a Inginerilor si Techicienilor din Romania)

Vol. 2, no. 5, Sept/Oct. 1958  
Bucuresti, Rumania

Monthly list of European Accession Index (EEAI) LC Vol. 8, no. 11  
November 1959  
Uncl.

TUDOROIU, V.

Automation in the extraction, transportation, and distribution of  
methane gas. p. 104.

AUTOMATICA SI ELECTRONICA. (Asociatia Stiinfitica a Inginerilor si  
Tehnicienilor din Romania)  
Bucuresti, Rumania  
Vol. 3, no. 3, May/June 1959.

Monthly list of Eastern European Accession Index (EEAI) LC Vol. 8, No. 11  
November 1959  
Uncl.

TUDOROV, Iv.

Treatment of false prognathism and of overbite with the apparatus  
fixed directly in the mouth. Stomatologija, Sofia no.5:313-315 1954.

1. Is okrushnata stomatologicheska poliklinika, Sofiia.  
(MALOCCLUSION, therapy,  
appar., fixation directly in mouth)

TUDOROVSKAYA, G.L.; MARGOLIS, F.G.

Kinetics of urea decomposition in the presence of monoammonium phosphate. Trudy NIUIF no.208:3-7 '65.

Quaternary system of urea, ammonium nitrate, monoammonium phosphate at 25°C. Ibid.:7-16 (MIRA 18:11)

B  
 SINGULARITIES IN THE VARIATIONS OF THE REFRACTIVE INDEX OF GLASS BELOW 300°. N. Tugorovskaja (Compt. rend. Acad. Sci. U.R.S.S., 1936, 1, 27-30).-- With Na<sub>2</sub>O-SiO<sub>2</sub> glass below the annealing temp. n increases with temp., but, on cooling, vals. of n > before are obtained, except at low temp. At const. temp. n increases with time, becoming const. in approx. 10-15 min. On Heating Na<sub>2</sub>O glass, quenching in glycerol, and comparing with unheated glass, n decreases in 2-3 min. after placing the glass in the furnace at < 170-200°, and then becomes const. Above 170-200° n decreases tp a ,on/ on 2-3 min., the change being > before, and then rises to a const. val. The final change in n is the same in both cases, and the time of the change is independent of temp.

R. S. B.

B-1-9

AIR-108 METALLURGICAL LITERATURE CLASSIFICATION

STANDARD INDEX  
CLASSIFICATION  
NUMBER

*Sa**A 53  
i*

1648. Variation of Refractive Index of Glass at Temperatures Below 300°. N. Tadevosyan. Compte Rendus (Doklady) de l'Acad. des Sciences, U.S.S.R., No. 117, p. 87-90, 1958. In French.—Glasses of composition  $\text{Na}_2\text{O}$ ,  $\text{SiO}_2$ ,  $\text{PbO}$ ,  $\text{SiO}_3$  and a barium crown are tested to find the alteration in their refractive indices after heating. It is found that for constant temperature there is an increase of refractive index. It is found that for of the increases or decrease during heating. Also, the time taken for the refractive index to become constant does not depend appreciably on temperature or composition of the glasses but on the size of the change in refractive index in comparison with the final value. H. M. B.

AB-34 METALLURGICAL LITERATURE CLASSIFICATION

SUBJ. INDEX	TOPIC INDEX	100000 AND OVER ONE	VALUATION	EIGHT BULLETIN											
				1	2	3	4	5	6	7	8	9	10	11	12


Some singularities in the variations of the index of refraction of glass at temperatures below 300° N. Tukrovskaya, Compt. rend. acad. sci. U. R. S. S. [N. S.] 1, 27-30 (1936).--When glass was quickly heated to temps. below the annealing temp. and then kept at a const. temp.,  $n$  increased to a const. value during 10-15 min. regardless of whether  $n$  had increased or decreased while the glass was being heated. The cooling curve was always above the heating curve but approached it as the temp. dropped. The time required for  $n$  to become const. depended only on the amt. that  $n$  varied and was independent of the temp. or the compn. of the glass. Simple Na<sub>2</sub>O-SiO<sub>2</sub> and PbO-SiO<sub>2</sub> glasses and a Ba crown glass all showed the same phenomenon. A glass contg. 77% SiO<sub>2</sub> and 23% Na<sub>2</sub>O was used in the following expts. Test pieces were placed in a const.-temp. oven for 1, 1.5, 2, etc., min. and then tempered by being quenched in glycerol at room temp. The  $n$  was compared with its value before

tempering. At oven temps. of 120-170°  $n$  dropped in about 4 min. to a const. value which was independent of the oven temp. At oven temps. above 170°  $n$  dropped to a min. in about 2.5 min. and then rose to the same const. value as was obtained below 170°. At temps. of 80-120°  $n$  dropped to values which were const. at any given temp. but which dropped as the temp. increased. From 170° to 200° the value of the min. dropped; above 200° the min.  $n$  was const. When plotted against temp. the const. values between 80° and 120° and the mins. between 170° and 200° fell on a straight line. The values of  $n$  in these tempered pieces remained const. for at least 3 months but returned to its normal value when they were annealed at 300°.

John E. Milbery

19

A.S.T.M. METALLURGICAL LITERATURE CLASSIFICATION



"APPROVED FOR RELEASE: 03/14/2001

CIA-RDP86-00513R001757410002-4

TUDOROVSKAYA, N.A.; SHERSTYUK, A.I.

Studying the process of catalyzed crystallization by the method of  
differential thermal analysis. Stekloobr. sost. no.1:119-122 '63,  
(MIRA 17:10)

APPROVED FOR RELEASE: 03/14/2001

CIA-RDP86-00513R001757410002-4"

ALEKSEYEV, A.G.; VARGIN, V.V.; VERTSNER, V.N.; KIND, N.Ye.;  
KONDRAT'YEV, Yu.N.; PODUSHKO, Ye.V.; SEREBRYAKOVA, M.V.;  
TIKHOMIROV, G.P.; TUDOROVSKAYA, N.A.; FLORINSKAYA, V.A.;  
LIBERMAN, N.R., red.

[Controlled catalyzed crystallization of glasses of the  
lithium aluminosilicate system] Katalizirovannaia regu-  
liruemaya kristallizatsiia stekol litievoalumosilikatnoi  
sistemy. Leningrad, Khimiia. Pt.1. 1964. 119 p.  
(MIRA 18:4)

ACCESSION NR: AT4019300

S/0000/63/003/001/0119/0122

AUTHOR: Tudorovskaya, N. A. ; Sherstyuk, A. I.

TITLE: Study of the process of catalyzed crystallization by the method of differential thermal analysis

SOURCE: Simpozium po stekloobraznomu sostoyaniyu. Leningrad, 1962. Stekloobraznoye sostoyaniye, vy\*p. 1: Katalizirovannaya kristallizatsiya stekla (Vitreous state, no. 1: Catalyzing crystallization of glass). Trudy\* simpoziuma, v. 3, no. 1. Moscow, Izd-vo AN SSSR, 1963, 119-122

TOPIC TAGS: thermal analysis, glass, glass crystallization, petalite, spodumene, catalyzed crystallization, titanium dioxide, alumina silicate

ABSTRACT: The thermal effect of crystallization was investigated in glasses of the system  $\text{Li}_2\text{O}-\text{Al}_2\text{O}_3-\text{SiO}_2$  having the composition of petalite or spodumene with admixtures of  $\text{TiO}_2$  and other oxides in amounts less than 10% by weight. X-ray and mineralogical analysis showed that in glass having a composition close to spodumene, the first thermal effect is produced by the crystallization of the high-temperature spodumene. The presence of the second high-temperature effect shows that a second, more refractory

Card 1/2

ACCESSION NR: AT4019300

phase is also crystallized in these glasses. For the two glasses investigated (glass 13 with 5% TiO<sub>2</sub> and spodumene glass), thermograms were obtained at a heating rate of 2-29 degrees/minute. The time during which the thermal effect disappears completely at a given temperature of exposure determine the rate of crystallization of the glass at this temperature. For glass 13, the effect of TiO<sub>2</sub> (1-11% by weight) on the thermal effect was also investigated, thermograms being plotted at a heating rate of 7 degrees/min. On the basis of the amount and character of the exothermal effects on the thermogram, the amount of each crystallized phase could be determined. Surface crystallization and overall crystallization could also be distinguished on the thermograms. The relative amount of crystallized phase was calculated from the thermograms by the method fo A. G. Vlasov and A. I. Sherstyuk. Orig. art. has: 2 figures and 2 tables.

ASSOCIATION: None

SUBMITTED: 17May63

DATE ACQ: 21Nov63

ENCL: 00

SUB CODE: MT

NO REF SOV: 001

OTHER: 000

Card 2/2

TUDOROVSKAYA, N. A.

TUDOROVSKAYA, N. A.: "Structural changes in the indexes of refraction of glass at temperatures below 300 degrees Centigrade." State Order of Lenin Optical Inst imeni S. I. Vavilov. Len-ingrad, 1956. (Dissertation for the Degree of Candidate in Physicomathematical Sciences).

SO: Knizhaya Istotopis', No 23, 1956

SELEZNEVA, Yevgeniya Semenovna; TUDOROVSKAYA, Yelena Aleksandrovna;  
KLADO, T.N., otv.red.; SOLOVEYCHIK, A.A., tekhn.red.

[P.A.Molchanov; eminent Soviet aerologist] P.A.Molchanov  
vydaiushchiisia sovetskii aerolog. Leningrad, Gidrometeor.  
izd-vo, 1958. 101 p. (MIRA 12:2)  
(Molchanov, Pavel Aleksandrovich, 1893-1941)  
(Meteorology)

MAYOROV, S.A.; YEVTEYEV, F.Ye., prof., retsenzent; TUDOROVSKIY,  
A.A., kand. tekhn. nauk, red.

[Technology of the manufacture of computers] Tekhnologija  
proizvodstva vychislitel'nykh mashin. Moskva, Mashino-  
stroenie, 1965. 410 p. (MIRA 18:9)

TUDOR ALEXANDER  
ILLARIONOVICH

'764

Optics  
optical instruments  
illumination

1875-1963

KLIMOV, V.P.; ROZEMBERG, Ya.G.; TUDOROVSKIY, V.P., otv.red.; NOVIKOVA, Ye.S., red.; KARABILLOVA, S.F., tekhn.red.

[Suggestions of efficiency experts on wire-broadcasting networks and electric communications within districts] Ratsionalizatorskie predlozheniia po vnutriraionnoi elektrorviazi i radiofikatsii. Moskva, Gos.isd-vo lit-ry po voprosam sviasi i radio, 1960. 55 p. (MIRA 14:3)

(Wire broadcasting) (Telecommunication)

TULOREKIY, Yu.

Lozh i litsemeriye Amerikanskoy burzhuaznoy demokratii (Lie's and hypocrisy of American bourgeois democracy, by) V. Ivanov i Yu. Tulorekiy. Moskva, Gospolitizdat, 1951. 122 p. Catalogued from abstract. Discusses the poverty and starvation facing the American people, the American workers' lack of rights, the fascistic trend of life in the USA, etc.

V/8  
850  
.19

"APPROVED FOR RELEASE: 03/14/2001

CIA-RDP86-00513R001757410002-4

Photochemical reaction of the isomers of carbon monoxide.

Isomerization of the isomers of carbon monoxide.

APPROVED FOR RELEASE: 03/14/2001

CIA-RDP86-00513R001757410002-4"

valid, also for the quaternary ammonium cations, such as the Cl-catalyzed decoupling of  $\text{O}_2$ . The following scheme (C.A. 45, 8394g) was used as the basis of calculation:  $\text{O}_2 + \text{Cl} \rightarrow \text{ClO} + \text{O}_2$ ,  $\text{ClO} + \text{O}_2 \rightarrow \text{Cl} + \text{O}_3$ ,  $\text{O}_3 + \text{O}_2 \rightarrow 2\text{O}_2$ .

(2)

"APPROVED FOR RELEASE: 03/14/2001

CIA-RDP86-00513R001757410002-4

~~ALL INFORMATION CONTAINED  
HEREIN IS UNCLASSIFIED~~

APPROVED FOR RELEASE: 03/14/2001

CIA-RDP86-00513R001757410002-4"

HUSS, F., Central Chemical Research Institute, Hungarian A.S. Budapest

"Study of the Inhibition of the Thermal Polymerization of Styrene," a paper submitted at the International Symposium on Macromolecular Chemistry, 9-15 Sep 1957, Prague.

Distr: bE2c(j)

7  
2-17-74  
Kinetics of the inhibition of the thermal polymerization of styrene! I. Kinetics of the one-step inhibition. F. Tudos and N. I. Smirnov (Hungarian Acad. Sci., Budapest, post). *Acta Chim. Acad. Sci. Hung.*, 15, 389-99 (1938); (in German).—The kinetics of the 1-step inhibition of polymerization through free radicals was investigated, and equations for the concn. of active centers established. II. Mechanism of the two-step inhibition. *Ibid.* 401-8.—The kinetics of the consecutive reactions between inhibitor, and free radicals was developed. III. Kinetics of the copolymerization of the inhibitor. *Ibid.* 409-15.—Equations were derived to investigate the copolymerization of inhibitors. The increase of the length of the inhibition period as a function of the concn. of inhibitor was not linear. IV. Thermal polymerization of styrene inhibited by quinones. F. Tudos and V. Kuri. *Ibid.* 417-39.—Benzquinone inhibited the 2-step polymerization of styrene. A semiquinoidal radical formed as an intermediate. Activation energies of some elementary reactions were detd. V. Mechanism of the effect of stable free radicals. *Ibid.* 541-7.—It was shown that 1,1-diphenyl-2-picrylhydrazyl (cf. Poirier, et al. *C.A.*, 47, 8071b), in addn. to inhibiting, also simultaneously initiates the polymerisation of styrene. *Milford Malenthal*

TUDOS, F.

Distr: 4E2c(j) 7  
Inhibited thermal polymerization of styrene; kinetics of copolymerization of quinones with styrene. F. Tudos (Hungarian Acad. Sci., Budapest). *J. Polymer Sci.* 30, 333-50 (1958).—A plot of conversion vs. time shows an inflection point if concn. of inhibitor is higher than a certain value. The curve of initial polymerization velocity against concn. of inhibitor has a min. at a certain concn. Expts. with *p*-benzoquinone at 90, 105, 120, and 135° agree with these conclusions. Activation energy of the inhibition reaction is 4.3 kcal./mol., and the activation energy of copolymerization of quinone is 9.4 kcal./mol. M. H. Danzig

JW  
1/1

4  
2 MAY  
1

gj

TUDOS, F.

SCIENCE

PERIODICALS: ACTA ZOOLOGICA. Vol. 64, No. 7/8 July/Aug. 1958  
MACYAR KEMIAI FOLYOTRAT

Tudos, F. Heat-transfer problems of polymerization-kinetic investigations.  
p. 270

Monthly list of East European Accessions (EEAI) LC, VOL. 8, No. 2,  
February 1959, Unclass.

TUDOS, F.

SCIENCE

PERIODICAL: MAGYAR KEMIAI FOLYOIRAT. Vol. 64, no. 7/8, July/Aug. 1958

Tudos, F. Investigation into some stabile free radicals. p. 305.

Monthly list of East European Accessions (EEAI) LC, Vol. 8, No. 2,  
February 1959, Unclass.

TUDOS, Ferenc, a kemiai tudomanyok kandidatusa (Budapest)

Kinetics of polymerization processes. I. Kem.tud.kozl.MTA 12 no.4:  
455-471 '59. (MAI 9:4)

1. A Magyar Tudomanyos Akademia Kozponti Kemiai Kutato Intezete,  
Budapest.

(Polymers and polymerization)

TUDOS, F.

## PAGE I BOOK INFORMATION

SOY/983

*International symposium po makromolekularej chemicz. Minsk, 12-15 lipca 1960. Księga 1 i 2. Sekretarz T. Sieratzy. II. (International Symposium on Macromolecular Chemistry Held in Minsk, June 12-15, Papers and Summaries). Section II. [Minsk, Izd-vo M. SSSR, 1960]. 559 p., 55,000 copies printed.*

Sponsoring Agency: The International Union of Pure and Applied Chemistry, Committee on Macromolecular Chemistry

Tech. Rep.: T.A. Presakov.

PURPOSE: This book is intended for chemists interested in polymerization reactions and the synthesis of macromolecular compounds.

CONTENTS: This is Section II of a multivolume work containing papers on macromolecular chemistry. The papers in this volume treat mainly the kinetics of thermal polymerisation reactions initiated by different catalysts or induced by radiation. Among the research techniques discussed are electron paramagnetic resonance spectroscopy and light-scattering intercalation. There are summaries in English, French and Russian. No page numbers are mentioned. References follow each article.

Bogdan'yan, E.I., and Z.I. Shustova. (rus.). Inhibition of Polymerisation by Aromatic Compounds 22

Gidde, Z., J. Konda, and M. Arai. (Hungary). Kinetics of the Inhibition of Polymerisation of Styrene by Nitro Compounds 31

Kazmierczak, G.A., L.M. Pernicki, V.R. Litchfield, and V.J. Killa. (USA). Radical Decomposition Reactions of Some Ferrocyanides and Pyroferrocyanides 53

Elshabani, J.A., and O.A. Flisovskiy. (USSR). On the Relative Activity of Butadiene-1,3-pentadiene in Polymerisation and Co-polymerisation Reactions With Other Dienic Compounds 62

Furukawa, J.M., and S.Y. Frechell. (USA). Intramolecular Exchange Reactions in the Process of Radical Polymerisation 72

Hajdu, D., L. Jánosi, G. Károly, and V.P. Li. (Hungary). Kinetic Study of Radical Polymerisation of Vinyl Monomers in the Presence of  $\text{SiCl}_4$  103

Krivchenko, M., and B. Grodzinska. (Poland). A Method of Measuring the Polymerisation Rate at a High Degree of Conversion 120

Kubisa, J., and M.F. Kiseleva. (rus.). Study of the Mechanism of Radiation Polymerisation 127

Lukáček, A., and M. Šloufek. (Czechoslovakia). The Polymerisation Rate for a Single Particle During Emulsion Polymerisation 135

Frants, P., and Ya. Sabirov. (Czechoslovakia). Emulsion Polymerisation of Chitosane 149

Purush, P., and G. Weissman. (Poland). Change of Potential During Polymerisation in Carbonyl-Substituted Systems 157

Radíček, Z., and A. Kefřík. (Czechoslovakia). The Heat of Reaction As a Means of Studying the Mechanism of the Emulsion Polymerisation of Styrene and Chitosane 165

Sedlák, Fr., D.F. Polozikov, A.N. Gerasimov, and S.I. Melnikov. (USSR). Polymerisation in the Presence of Organic Compounds or Alkali Metals 169

Korobov, A.B., D.P. Matveeva, and V.B. Kravtsova. (USSR). On the Polymerisation Kinetics and Mechanism of the Polymerisation of Methyl Methacrylate by Polyvinylchloride 203

Dulera, M., M. Jelínek, I. Lančová, and K. Veselý. (Czechoslovakia). Chain Depolymerisation During the Anionic Polymerisation of Octaethylcyclotetrasiloxane. The Formation of Stable Complexes as Active Centers 210

Michálek, Z., I. Michál, and J. Plíš. (Czechoslovakia). Kinetics of the Polymerisation of Formaldehyde 213

Veselý, E. (Czechoslovakia). On the Mechanism of Ionic Polymerisation 262

Kralíček, Z., and A. Kefřík. (Czechoslovakia). On the Role of Nonpolar Compounds in the Cationic Polymerisation of Isobutylene 272

45

TUDOS, Ferenc (Budapest XIV Hungaria korut 114); BEREZHNIKH, Tamara F.  
(Budapest XIV Hungaria korut 114); AZORI, Maria (Budapest XIV  
Hungaria korut 114)

Kinetics of the inhibition of the polymerization of styrol. I. Effect  
of the stable-free radicals. Acta chimica Hung 24 no.1:91-106 '60.  
(EEAI-10:4)

1. Central Research Institute for Chemistry, Hungarian Academy of  
Sciences, Budapest.

(Polymers and polymerization) (Styrene)  
(Diphenylpicrylhydrazyl)

IRVING, Rezsone; TUDOS, Ferenc; TELEKI, Piroska

Initiated polymerization of styrol by sulfinic acid. I.  
Polymerization of styrol in the presence of p-toluolsulfonic acid as well as of benzoyl peroxide-p-toluolsulfonic acid system. Magy kem folyoir 66 no.10:415-422 O '60.

1. Kabel- es Muanyaggyar Kozponti Laboratorium, Budapest;  
Magyar Tudomanyos Akademia Kozponti Keriai Kutato Intezete,  
Budapest; es Budapesti Muszaki Egyetem Muanyag- es Gumiipari  
Tanszeke.

Hungary; 1950s; 1960s; 1970s

Committee of Central Control, Hungarian Socialist Republic (for 1950s)  
324-733-A2-1124

1. Central Research Institute of Chemistry of the Hungarian Academy of Sciences, Budapest (for 1950s and 1960s).  
2. Central Research Institute of Physics of the Hungarian Academy of Sciences, Budapest (for 1960s).

FOLDESNE BEREZSNICH, Tamara; TUDOS, Ferenc

Kinetics of radical polymerization. Pt.12. Magy kem folyoir  
70 no.11;500-504 N '64.

1. Central Research Institute of Chemistry, Hungarian Academy  
of Sciences, Budapest.

KIRALY, János (Budapest, XIV., Hungaria korut 114); FEJES, Pal (Budapest, XIV., Hungaria korut 114); TUDOS, Ferenc (Budapest, XIV., Hungaria korut 114); AZORI, Maria (Budapest, XIV., Hungaria korut 114)

Adsorption of oxygen on free radicals of 1,1-diphenyl-2-picryl hydrazyl.  
Acta chimica Hung 29 no.4:409-418 '61.

1. Central Research Institute for Chemistry, Hungarian Academy of Sciences, Budapest. 2. Editorial board member, "Acta Chimica Academiae Scientiarum Hungaricae" (for Tudos).

5,3200

39852  
S/190/62/004/008/014/016  
B101/B138

AUTHORS: Tüdös, F., Kende, I., Azori, M.

TITLE: Kinetics of inhibition of radical polymerization. IV. Effect of mono- and dinitro-benzene derivatives on the induced polymerization of styrene

PERIODICAL: Vysokomolekulyarnye soyedineniya, v.4, no. 8, 1962, 1262-1270

TEXT: The influence of substituents on the inhibiting effect of nitro-aryl compounds was studied during the polymerization of styrene induced by azoisobutyric acid dinitrile, and also the dependence of chain regeneration on polar factors. Modified kinetic equations including those by L. J. Kice (J. Amer. Chem. Soc., 76, 6274, 1954) and data obtained by P. D. Bartlett, H. Kwart (J. Amer. Chem. Soc., 72, 1051, 1950; ibid., 74, 3969, 1952), D. H. McDaniel and H. C. Brown (J. Organ. Chem., 23, 420, 1958) were used to calculate the reactivity  $k_5/k_2$ .the reactivity nitro group  $\frac{1}{n}k_5/k_2$ , and the algebraic sum  $\sum \sigma_i$  of the

Card 1/2

S/190/62/004/008/014/016  
B101/B138

Kinetics of inhibition of radical ...

Hammett constants of substituents, for nitro-benzene, o- and p-nitro-toluene, p-nitro-chloro benzene, p-nitro-anisol, ethyl ester of p-nitrobenzoic acid, o-, m-, and p-dinitro-benzene, 2,4-dinitro-toluene (Table). Results: The effect of the substituent can be described satisfactorily by Hammett's equation. Electron acceptors increase relative reactivity and reduce chain propagation. The reaction of nitro compounds with polystyrene is more sensitive to the effect of polar factors than that with methyl acrylate or vinyl acetate. This is due to compensation of the effect of electron acceptor groups supporting the latter reaction, by electrostatic repulsion between the m and nitro groups. There are 3 figures and 1 table.

ASSOCIATION: Tsentral'nyy nauchno-issledovatel'skiy institut khimii  
AN Vengrii; Budapesht  
(Central Scientific Research Institute of Chemistry of the  
AS Hungary, Budapest)

SUBMITTED: December 28, 1961

Card 2/2

5.3832

5.3200

AUTHORS: Tüdös, F., Simandi, L.

TITLE: Kinetics of inhibition of radical polymerization. V. Effect of p-benzoquinone and its derivatives on the induced polymerization of styrene

PERIODICAL: Vysokomolekulyarnyye soyedineniya, v. 4, no. 8, 1962;  
1271-1281

TEXT: The polymerization of styrene induced by azoisobutyric acid dinitrile was conducted in the presence of benzoquinone, toluquinone, o-xyloquinone, p-xyloquinone, trimethyl-p-benzoquinone, methoxyquinone, or monochloro-n-benzoquinone at 50°C by methods described previously (MTA KKKI Közleményei, no. 3, 3, 1959; ibid., no. 5, 13, 1961; Acta Chim. Hung., 14, 417, 1958). The degree of conversion was determined dilatometrically. The stoichiometric coefficient  $\mu$  and the relative reactivity  $k_5/k_2$  of the inhibitors were calculated from equations derived previously (MTA KKKI, Közleményei, no. 2, 51, 1959) (Table 2). Results:

Card 1/3

39853  
S/190/62/004/008/015/016  
B101/B180

S/190/62/004/008/015/016  
B101/B190

Kinetics of inhibition of radical ...

There is a strictly linear dependence between  $\log k_5/k_2$  and the redox potential  $E^{\circ}$ . The data for  $E^{\circ}$  were taken from publications by D. E. Kvalnes (J. Amer. Chem. Soc., 56, 667, 670, 1934) and W. M. Clark (Oxidation-Reduction Potentials of Organic Systems. The Williams and Wilkins Co., Baltimore, 1960, 371). The slope of the straight line is  $\alpha F/RT$ , where  $\alpha = 1/2$ . Hence, this reaction is ideally symmetrical according to O. Dimroth (Angew. Chemie, 46, 571, 1933). The stoichiometric coefficient  $\mu$  is lower than the theoretical value. It decreases as the number of substituents increases, its value depending on the position of the latter. Hence, part of the radicals formed from the inhibitor are assumed to react directly with the monomer and cause partial copolymerization of quinone and the monomer.  $2/\mu$  inhibitor molecules join in with the chain which is in good agreement with data obtained by J. C. Bevington et al. (J. Chem. Soc., 1955, 2822). There are 4 figures and 2 tables.

ASSOCIATION: Tsentral'nyy nauchno-issledovatel'skiy institut khimii AN Vengrii, Budapest (Central Scientific Research Institute of Chemistry of the AS Hungary, Budapest)

Card 2/3

Kinetics of inhibition of radical ...

S/190/62/004/008/015/016  
B101/B180

SUBMITTED: December 28, 1961

Table 2. Kinetic data of the inhibitors studied. Legend: (1) inhibitor;  
 (2) benzoquinone; (3) toluquinone; (4) methoxyquinone; (5) p-xyloquinone;  
 (6) o-xyloquinone; (7) monochloro-p-benzoquinone; (8) trimethyl-p-  
 benzoquinone.

Ингибитор	$\mu$	$k_1/k_2$	$F^a$
BX 2	1,26	518±25	0,711
TX 3	1,48	266±15	0,653
MOX 4	1,08	103±10	0,642
n-KX 5	1,05	82±10	0,604
o-KX 6	0,70	120±20	
XX 7	1,10	720±70	0,734
TMX 8	0,85	25±5	0,527

Card 3/3

S/190/62/004/009/012/014  
B101/B144

54300.

AUTHORS: Tüdös, F., Simándi, L.

TITLE: Inhibition kinetics of radical polymerization. VI. Effect of p-xyloquinone on the initiated polymerization of styrene

PERIODICAL: Vysokomolekulyarnyye soyedineniya, v. 4, no. 9, 1962, 1425-1430

TEXT: The inhibiting effect of p-xyloquinone on the initiated polymerization of xylene at 40 and 60°C was studied in the same way as previously (MTA KKKI Közleményei, no. 5, 13, 29, 1961; ibid., no. 7). Results: (1) The relative reactivity  $k_5/k_2$  is a linear function of  $1/T$ .  $k_5/k_2$  is 106 at 40°C and 61 at 60°C. (2) The activation energy of the inhibition is zero, the frequency factor is very low:  $\log A_5 = 5.04 \pm 0.31$ . (3) Hence, inhibition is a nonadiabatic and exothermic reaction with a very low transmission coefficient; an approximate calculation for p-benzoquinone gave  $\Delta H \geq -10$  kcal/mole. (4) The stoichiometric coefficient, which is 1.06 at 40°C and 0.97 at 60°C, is independent of the temperature. There are 1 figure and 2 tables.

Card 1/2

JC

Inhibition kinetics of...

S/190/62/004/009/012/014  
B101/B144

ASSOCIATION: Central Scientific Research Institute of Chemistry,  
Hungarian AS, Budapest

SUBMITTED: December 28, 1961

Jc

Card 2/2

5.4300

S/190/62/004/009/013/014  
B101/B144AUTHORS: Tüdös, F., Simándi, L., Azori, M.

TITLE: Inhibition kinetics of radical polymerization. VII. Effect of halogenated quinones on the initiated polymerization of styrene

PERIODICAL: Vysokomolekulyarnyye soyedineniya, v. 4, no. 9, 1962, 1431-1444

TEXT: Inhibition of the polymerization of styrene by 2,5-dichloro benzoquinone, 2,6-dichloro benzoquinone, trichloro benzoquinone, chloranil, bromanil, and iordanil at 50°C was studied by methods described previously (MTA KKKI Közleményei, no. 7 (V és VI); ibid. no. 5, '61, 1961). Halogenated quinones were found to be active inhibitors. The inhibition period, however, is a nonlinear function of the inhibitor concentration. A radical which causes chain regeneration is formed by halogenated quinones with the monomer at a ratio of 1:1. The formation of this "charge transfer" complex (R. S. Mulliken, J. Amer. Chem. Soc., 74, 811, 1952) increases the reactivity of the halogenated quinones owing to an increase in transmission coefficient. The stoichiometric coefficient is

Card 1/2

JC

Inhibition kinetics of...

S/190/62/004/009/013/014  
B101/B144

VC

a nontrivial quantity and decreases as the number and size of substituents increase. On the basis of the implicit equation  $I(z) = \text{const} - w_1 ct / 4\mu$  for the dependence of the inhibitor concentration on time, which had been developed previously (J. Polymer Sci., 30, 343, 1958), the equations  $t_{i,r} = (4\mu / 2k_1 f c') I(z_0/x_0)$  and  $t_{i,r} = t_i (4x_0/c' z_0) I(z_0, x_0) \equiv t_i I^*(z_0/x_0)$  (without chain regeneration), where  $c' = cx_0$ , are derived for the inhibition period. The experimental data are in good agreement with the theoretical values. There are 6 figures and 8 tables.

ASSOCIATION: Central Scientific Research Institute of Chemistry,  
Hungarian AS, Budapest

SUBMITTED: December 28, 1961

Card 2/2

SZPICIN, V.I. [Spitsin, V.I.]; TUDOS, Ferenc [translator]

Present state of chemical sciences and their long-range development in the Soviet Union. Kem tud kozl MTA 18 no.2:173-219 '62.

1. A. Szovjetunio Tudomanyos Akademaja Fizikai Kemial Intezete, Moscow, es Lomonoszov Egyetem, Moscow (for Spitsin). 2. "A Magyar Tudomanyos Akademia Kemial Tudomanyok Osztalyanak Kozlemenyei" szerkeszto bizottsagi tagja (for Tudos).

SZPICIN, V.I. [Spitsin, V.I.]; TUDOS, Ferenc [translator]

New data on the effect of the radioactive radiation of solid bodies, as well as of external radiation on certain heterogeneous chemical processes. Kem tud kozl MTA 18 no.2: 301-321 '62.

1. Szovjetunio Tudomanyos Akademaja Fizikai-kemial Intezete, Moscow, es Lomonoszov Egyetem, Moscow (for Spitsin). 2. "A Magyar Tudomanyos Akademia Kemial Tudomanyok Osztalyanak Kozlemenyei" szerkeszto bizottsagi tagja (for Tudos).

ERDEY-CRUZ, Tibor, akademikus; BRUCKNER, Gyozo, akademikus; LENGYEL, Bela; TELEGY-D-KOVATS, Laszlo, a tudomanyok doktora; HARDY, Gyula, kandidatus; GERECS, Arpad, akademikus; FOLDI, Zoltan; WOLKÖBER, Zoltan; TUDOS, Ferenc, kandidatus; PURMAN, Jeno; KRAUSZ, Imre, kandidatus; ERDEY, Laszlo, akademikus; SCHAY, Geza, akademikus

An account of the 1961 work of the Section of Chemical Sciences, Hungarian Academy of Sciences. Kem tud kozl 18 no.3:343-394 '62.

1. Magyar Tudomanyos Akademia Kemial Tudomanyok Osztalyanak titkara, es "A Magyar Tudomanyos Akademia Kemial Tudomanyok Osztalyanak Kozlemenyei" szerkesztoje (for Erdey-Gruz). 2. Akademial levelezo tag (for Lengyel and Foldi), 3. "A Magyar Tudomanyos Akademia Kemial Tudomanyok Osztalyank Kozlemenyei" szerkeszto bizottsagi tagja (for Bruckner, Erdey, Foldi, Gerecs, Hardy, Lengyel, Schay, Tudos).

HARDY, Gyulane; RADICS, Lajos; TUDOS, Ferenc, kandidatus

An account of the 5th International Symposium on Free Radicals.  
Kem tud kozl. 18 no.3:509-519 '62.

1. Magyar Tudomanyos Akademia Kozponti Kemial Kutato Intezete,  
Budapest. 2. "A Magyar Tudomanyos Akademia Kemial Tudomanyok  
Osztalyanak Kozlemenyei" szerkeszto bizottsagi tagja (for Tudos).

TUDOS, Ferenc, a kémiai tudományok Kandidátusa

Discussion of the kinetics of radical polymerization on the basis of the hypothesis of hot radicals. J. Polym. Sci. A-1 1964, MFA 21 no. 1849-87 '64.

J. Central Research Institute of Chemistry, Hungarian Academy of Sciences, Budapest; Editorial board member, "A Magyar Tudományos Akadémia Kemiai Tudományok Osztályának Folyóiratai."

33407

H/005/62/000/002/001/001  
D283/D304

15-8110

AUTHORS: Iring, Rezsö and Tüdös, Ferenc

TITLE: Polymerization of styrene initiated with sulfinic acid.  
II. Polymerization in the presence of a system of benzoyl-  
peroxide-intermetallic compound of different valencies-  
p-toluene sulfinic acidPERIODICAL: Magyar kémiai folyóirat<sup>68</sup>, no. 2, 1962, 86-92

TEXT: After a brief review of data contained in the first article of the series, the authors describe their experiments with polymerization of styrene carried out to determine the suitability of p-toluene sulfinic acid as a reducing agent in a system of benzoyl-peroxide and an intermetallic compound of different valencies. Data contained in the article have been taken from Mrs. Iring's dissertation for the degree of Candidate. The authors examined the polymerization of styrene with benzoyl-peroxide + copper acetyl acetonate + p-toluene sulfinic acid and with benzoyl-peroxide + iron acetyl acetonate + p-toluene sulfinic acid. The experiments proved

Card 1/3

33407

H/005/62/000/002/001/001

D283/D304

## Polymerization of styrene ...

that polymerization when initiated with the benzoyl-peroxide + copper-acetyl acetonate + p-toluene sulfinic acid does not depend on the concentration of p-toluene sulfinic acid. In case of variable p-toluene sulfinic acid concentration, the kinetic curves show the typical form of the "dead end" polymerization after a constant initial velocity, while the limit of conversion ( $x_{\infty}$ ) is in direct ratio to the square root of the p-toluene sulfinic acid concentration. The square of the initial velocity of polymerization is also in direct ratio to the copper acetyl acetonate concentration. The experimental results also proved that while the polymerization velocity increases with an increase of catalyst concentration, the limit of the obtainable final conversion decreases at the same time.

According to

$$\log(x_{\infty} - x) = \log x_{\infty} - \frac{k''}{4.6} t \quad (8)$$

the value of  $\log(x_{\infty} - x)$  decreases linearly with the time. The directional tangent of the set of curves is in direct ratio to the initial concentration of the catalyst.

$$\text{tg } \alpha = \frac{k'' i}{4.6} = \frac{k_i \text{ [copper acetyl acetonate]}}{4.6} \quad (9) \text{ which indicates that}$$

Card 2/3

33407

H/005/62/000/002/001/001  
D283/D304

Polymerization of styrene ...

the process of initiation is bimolecular. In case of the benzoyl peroxide + iron acetyl acetonate + p-toluene sulfinic acid system the increase of polymerization velocity is noticeable which is not the case with the benzoyl-peroxide + copper acetyl acetonate + p-toluene sulfinic acid system. The polymerization velocity decreases with the increase of the iron acetyl acetonate concentration, while the molecular weight increases. The intermetallic compounds used in the experiments were supplied by the Központi kémiai kutató intézet polimerizációs kinetikai csoportja (Polymerization Kinetics Team of the Central Chemical Research Institute). There are 14 figures, 5 tables and 5 references. 2 Soviet-bloc, 1 non-Soviet-bloc and 2 unidentified. The reference to the English-language publication reads as follows: A.V.Tobolsky: J.Amer.Soc., 80, 5927, 1958. (Abstracter's note: 2 unidentified references refer to German-language publications; it was not possible to establish whether they are East or West German.)

ASSOCIATION: Kábel - és műanyaggyár, központi laboratórium (Cable and Synthetic Material Plant, Central Laboratory) (Iring); Magyar tudományos akadémia központi kémiai kutató intézete (Central Chemical Research Institute of the Hungarian Academy of Sciences) (Tüdös)

Card 3/3

1. 17234-66 EMP(j) RM

SOURCE CODE: HU/2502/65/043/001/0063/0072

ACQ NR: A16009211

AUTHOR: Turcsanyi, Bela-Turciani, B.; Tudos, Ferenc-Tyudesh, F. (Doctor)

52

FH

ORG: Central Research Institute for Chemistry, Hungarian Academy of Sciences,  
Budapest

TITLE: Organic molecular compounds. Part 2: Crystalline molecular compounds of  
pyrene with 1,4-benzoquinone and its substituted derivatives

SOURCE: Academia scientiarum hungaricae. Acta chimica, v. 43, no. 1, 1965, 63-72

TOPIC TAGS: IR spectroscopy, electron spin resonance, crystal chemistry, molecular  
crystal, quinone, crystal structure, chemical stability

ABSTRACT:

The crystalline molecular compounds of pyrene formed with 1,4-benzoquinone and its methyl and haloid substituted derivatives were prepared and characterized by their color, composition, and melting point. In respect to the selectivity of the formation of molecular compounds from quinones, the role of some factors affecting the stability of the crystalline complex was discussed. Some data on the infrared spectroscopical and electron spin resonance investigation of crystalline quinone-pyrene molecular compounds were presented. The authors express their thanks to the coworkers at the Institute, K. Shashvari and Sh. Kholl, for preliminary information prior to publication of the results of the radiological crystallographic and IR spectroscopic results. Further thanks is extended to the Faculty of Organic Chemistry at Szeged University for

2

Cord 1/2

L 17634-66

ACC NR: AT6009211

carrying out the analysis on molecular compounds on oxygen. Orig. art. has:  
4 figures and 2 tables. [JPRS]

SUB CODE: 20, 07 / SUBM DATE: 30Jul64 / ORIG REF: -004 / OTH REF: 0'6  
SOV REF: 001

EW  
Card 2/2

35812

H/005/62/000/004/001/001  
D249/D302

15.810<sup>0</sup>

AUTHORS:

Iring, Rezsöne, and Tüdös, Ferenc

TITLE:

Polymerization of styrene, initiated by sulphinic acid. III

PERIODICAL: Magyar kémiai folyóirat, no. 4, 1962, 149 - 151

TEXT: The effect of ethanol was investigated on the polymerization of styrene, initiated by the system p-toluene-sulphinic acid + benzoyl peroxide. It was found that a relatively small quantity of alcohol (3 - 5 %), which would not effect the rate of thermal or benzoyl peroxide, initiated polymerization and increased the rate of polymerization to a considerable extent. The curve indicating the rate of polymerization as a function of the concentration of alcohol passes through a maximum. This phenomenon is interpreted as due to the association of p-toluene-sulphinic acid when dissolved in alcohol. As a result of association the rate of monomolecular decomposition of p-toluene-sulphinic acid increases. Simultaneously the rate of reaction between p-toluene-sulphinic acid and benzoyl peroxide increases as well. The experimental data are presented.

Card 1/2

X

Polymerization of styrene, ...

H/005/62/000/004/001/001  
D249/D302

ted in tables and as kinetic curves. There are 3 figures, 2 tables and 3 references: 2 Soviet-bloc and 1 non-Soviet-bloc. The reference to the English-language publication reads as follows: W.G. Wright, J. Chem. Soc., 1919, 683.

ASSOCIATION: Budapest, kabél- és műanyaggyár (Factory of Cables and Plastics, Budapest)

SUBMITTED: August 3, 1961

Card 2/2

L 45346-66 EWP(j) JW/RM

ACC NR: AT6033616

SOURCE CODE: HU/2502/65/043/002/0245/0255

AUTHOR: Tudos, Ferenc--Tyudesh, F. (Doctor; Budapest); Heidt, Janos--Geidt, Y. (Budapest); Ero, J.--Ere, Y. (Budapest)

ORG: [Tudos; Heidt] Central Research Institute for Chemistry, MTA, Budapest (MTA Kozponti Kemial Kutato Intezet); [Ero] Central Research Institute for Physics, MTA, Budapest (MTA Kozponti Fizikai Kutato Intezet)

TITLE: Chemistry of free radicals. III. Synthesis and investigation of some stable free radicals of the Banfield type

SOURCE: Academia scientiarum hungaricae. Acta chimica, v. 43, no. 2, 1965, 245-255

TOPIC TAGS: free radical, condensation reaction, IR spectrum, electron spin resonance

ABSTRACT: A number of derivatives of some Banfield-type stable free radicals were synthesized with substitutions in the aromatic ring (o-, m- and p-CH<sub>3</sub>; m- and p-F; m- and p-Cl, m- and p-Br, and m-NO<sub>2</sub>). According to the experimental data, only such derivatives can be prepared in satisfactory yields which do not carry electron-donor or electron-acceptor substituents of overly excessive activity. The condensation reaction is, in general, impeded by substitution in the o-position. Experiments were also carried out to elucidate the mechanism of the condensation. The infrared spectra of the prepared compounds were recorded. In addition, the electron spin resonance spectra of the free radicals were also determined. Of the latter, the esr spectrum of the original compound and the interpretation of this spectrum are published. The authors thank Doctor A. Messmer and D. Chem for the valuable assistance in recording the infrared spectra, and Mrs. G. Szilagyi, Scientific officer for performing the microanalyses. Further thanks is extended to Miss J. Kozeluhu and Mr. L. Sumegi for their participation in the experimental work. Orig. art. has: 6 figures and 1 table. [Orig. art. in Eng.] [JPRS: 33,540]

SUB CODE: 07, 20 / SUBM DATE: 02Nov65 / ORIG REF: 003 / SOV REF: 001 / OTH REF: 013

Card 1/1

0720 1642

L 44609-66 EWP(j)/T RM

ACC NR: AT6033141

SOURCE CODE: HU/2502/65/044/004/0403/C457

AUTHOR: Tudos, F.—Tyuyesh, F.

ORG: Central Research Institute of Chemistry, MTA, Budapest (MTA Kozponti Kemial Kutatointezet) 35.  
34  
B+1

TITLE: Discussion of the kinetics of radical polymerization on the basis of the hypothesis of hot radicals II.

SOURCE: Academia scientiarum hungaricae. Acta chimica, v. 44, no. 4, 1965, 403-457

TOPIC TAGS: radical polymerization, polymerization kinetics

ABSTRACT: Experimental control of the hypothesis in case of inhibition: the value of the stoichiometric coefficient is independent of temperature; the effect of inert solvents; the synthesis of polymers which contain the inhibitor; dependence of the stoichiometric coefficient on the three dimensional structure of the inhibitor; dependence of the stoichiometric coefficient on other structural factors; dependence of the stoichiometric coefficient on the reactivity of the monomers. Generalization of the results. The effect of the reaction heat on the chain reactions; discussion of the rate of polymerization by postulation of the existence of hot radicals; discussion of the chain reaction by postulation of the existence of hot radicals; determination of the starting velocity with the help of the degrees of polymerization; discussion of other reactions by postulation of the existence of hot radicals; the

Card 1/2

L 44609-66

ACC NR: AT6033141

role played by the reaction heat in chain reactions. The author thanks the members of the Catalysis Committee, MTA, Professor G. Schay and Professor Z. Szabo for valuable discussions, advice and comments. He also thanks his colleagues in the Polymerization Kinetics Group, Central Research Institute for Chemistry, MTA, for their devoted cooperation and the research work which furnished the data published in this paper: Mr. I. Kende (nitro and nitroso compounds), Mrs. T. F. Bereznikh (aromatic hydrocarbons and vinyl monomers), and Mrs. T. L. Simandi (quinones, liquid-phase polymerizations). The author further thanks Mr. B. Turcsanyi and Mr. J. Heidt for preparing the inhibitors for the kinetic investigations, Mr. M. Simonyi for assistance in checking the data reported in the literature, and Mr. L. Sumegi and Mrs. K. B. Zsadon for their assistance. Orig. art. has: 8 figures, 97 formulas and 18 tables.  
[JPRS: 33,540]

SUB CODE: 07 / SUBM DATE: 15Sep64 / ORIG REF: 004 / SOV REF: 010  
OTH REF: 080

Card 2/2 blg

L 47422-66 EWP(j)/T RM

ACC NR: A76034995

SOURCE CODE: HU/0005/65/071/012/0524/0529

SIMONYI, Miklos, TUDOS, Ferenc, HEIDT, Janos, Hungarian Academy of Sciences,  
Central Research Institute of Chemistry (Magyar Tudomanyos Akademia, Kozponti  
Kemial Kutato Intezet), Budapest.

"Kinetics of Radical Polymerization XIII. Study of Metathetic Reactions by  
Means of an Inhibition Method." *25* *B*

Budapest, Magyar Kemial Folyoirat, Vol 71, No 12, Dec 65, pages 524-529.

TOPIC TAGS: radical polymerization, polymerization kinetics

Abstract: [Authors' English summary modified] An inhibition method was used to study the mechanism of some metathetic reactions. The reactivity of the inhibitors of this mechanism is generally slight; consequently, a new method has been developed to determine the length of the inhibition period and the reactivity of the inhibitor. The reactivity of the Banfield condensates studied showed a slight Hammett-dependency. The stoichiometric coefficients of metathetic reactions are also smaller than the theoretical ones, a fact looked upon as proof of the hot radical hypothesis. On the basis of the hypothesis, the Hammett-dependency was established for the reactivity of the intermediate radicals as well. A change of the other substituent on the nitrogen atom has a considerable influence on the kinetic parameters. Relative reactivity increases with the delocalization of the electron of the intermediate radical while a decrease in the space requirement of the substituent favors the regeneration of the hot radical. Orig. art. has: 4 figures, 15 formulas and 2 tables. [JPRS: 34,518]

SUB CODE: 07 / SUEM DATE: 20Apr65 / ORIG REF: 012 / SOV REF: 001 / OTH INF: 006  
Card. 1/1 vlr

0921 1506

L 46859-66 EWP(j)/T IJP(c) RM  
ACC NR: AP6034676

SOURCE CODE: HU/0005/66/000/005/0231/0237

411  
B

AUTHOR: Tudos, Ferenc--Tyudesh, F., Foldesne Bereznich, Tamara--Feldesne-Bereznykh, T.

ORG: Central Research Institute of Chemistry, Hungarian Academy of Sciences, Budapest  
(Magyar Tudomanyos Akademia, Kozponti Kemial Kutato Intezet)

TITLE: Kinetics of radical polymerization XVI. Condensed aromatic hydrocarbons II

SOURCE: Magyar kemial folyoirat, no. 5, 1966, 231-237

TOPIC TAGS: radical polymerization, aromatic hydrocarbon, activation energy

ABSTRACT: [Authors' Hungarian summary] The inhibition-kinetic study of some aromatic hydrocarbons is reported which exert a retarding effect on the polymerization of vinyl acetate. The dependence of reactivity on the parameters which characterize the structure of the aromatic hydrocarbon are discussed in detail. It was demonstrated that a radical attack can take place not only at the most reactive sites of the aromatic molecule but at less reactive sites as well, with correspondingly lesser probability. This circumstance was considered in the calculation of both reactivity and the theoretical value of the stoichiometric coefficient. The activation energy of the elemental inhibition reaction was discussed and it was shown that only a small fraction of the change in reactivity can be explained with the change in activation energy, the larger portion of it is related to a

Card 1/2

0921 1304

L 46859-66

ACC NR: AP6034676

Q

change in the pre-exponential factor. Therefore, the radical reactions of aromatic hydrocarbons can not be interpreted mechanically by using Polanyi's rule. Orig. art. has: 2 figures, 9 formulas, and 4 tables. [JPRS: 36,862]

SUB CODE: 07 / SUBM DATE: 17Sep65 / ORIG REF: 009 / SOV REF: 003  
OTH REF: 020.

Card 2/2

L 47239-66 EWP(j)/T RM

ACC NR: AI6034303

SOURCE CODE: HU/0005/66/000/006/0239/0243  
26  
BAUTHOR: Tudos, Ferenc; Ladik, Janos; Turcsanyi, BelaORG: Central Research Institute of Chemistry, Hungarian Academy of Sciences, Budapest  
(Magyar Tudomanyos Akademia, Kozponti Kemial Kutato Intezet)

TITLE: Kinetics of free radical polymerization XVII. Effect of charge transfer complexes on certain elemental processes of free radical polymerization

SOURCE: Magyar kemial folyoirat, no. 6, 1966, 239-243  
7

TOPIC TAGS: radical polymerization, polymerization kinetics, copolymerization

ABSTRACT: [Authors' English summary modified] Molecular compounds (charge transfer complexes) which are formed in some cases of radical polymerization have a considerable influence on the kinetics of the process. A theoretical study was made of the factors which determine the kinetic parameters of the reactions of radicals with other compounds which have a closed shell  $T^+$  electron system. Special attention was paid to the effect of the formation of molecular compounds. It was found that the increase in reactivity which is observed generally may be attributed to an increase in the resonance energy of the transition state. This can be used as a basis for the interpretation of some anomalous effects of inhibited polymerization and co-polymerization in a satisfactory manner. Orig. art. has: 3 figures and 17 formulas.  
[JPRS: 36,862]

SUB CODE: 07 / SUEM DATE: 19Jun65 / ORIG REF: 006 / SOV REF: 001  
OIH REF: 008  
Card 1/gd

092/ 0034

L 38647-66 EWP(j)/T RM

ACC NR: AP6027656

SOURCE CODE: HU/0005/66/000/004/0181/0186

AUTHOR: Foldesno, Berezsnich, Tamara--Feldeshne-Berezhnikh, T.; Tudos, Ferenc--  
Tyudash, F.ORG: Central Research Institute for Chemistry, MTA, Budapest (MTA Kozponti Kemial  
Kutato Intezete)TITLE: Kinetics of radical polymerization. Part 15: Investigation of condensed  
aromatic hydrocarbons. Section 1

SOURCE: Magyar kemial folyoirat, no. 4, 1966, 181-186

TOPIC TAGS: polymerization kinetics, radical polymerization, aromatic hydrocarbon,  
vinyl compound, acetate, stoichiometry

ABSTRACT: The inhibitory effects of perylene, 1,2,4,5-dibenzopyrene, 3,4-benzotetraphene, 1,2,3,4-dibenzopyrene, anthanthrene, 3,4-benzopyrene, 1,2-benzoanthracene, anthracene, tetracene, and 9-phenylanthracene in the radical polymerization of vinyl acetate were investigated at various temperatures. The effects of temperature were not very significant. The actual values for the stoichiometric coefficients were lower for all inhibitors than those calculated theoretically. The stoichiometric anomalies were interpreted in terms of hot-radical hypothesis. The authors thank J. Heidt and M. Vari for carrying out the experiments. Orig. art. has: 1 figure and 5 tables.

SUB CODE 1/1 07 / SUBM DATE: 17Sep65 / ORIG REF: 007 / SOV REF: 006 / OTH REF: 007

L 47238-66 EWP(J)/T RM

ACC NR: AP6034304

SOURCE CODE: HU/0005/66/000/006/0244/0248

AUTHOR: Simandi, Laszlone--Shimandine, L.; Tudos, Ferenc--Tyudesh, P.; Turcsanyi, Bela--Turchani, E.ORG: Central Research Institute of Chemistry, Hungarian Academy of Sciences, Budapest (Magyar Tudomanyos Akademia, Kozponti Kemial Kutato Intezet)28  
B

TITLE: Kinetics of free radical polymerization XVIII. Inhibition of free radical polymerization with quinone

SOURCE: Magyar kemial folyoirat, no. 6, 1966, 244-248

TOPIC TAGS: radical polymerization, polymerization kinetics, quinone, polymerization inhibition

ABSTRACT: [Authors' Hungarian summary] The inhibitory effect of multiple-ring quinones on the polymerization of styrol was examined. In addition to the member which responds to changes in the redox potential, the logarithm of relative reactivity of quinones also contains a member corresponding to the change in resonance energy which accompanies the transition from quinoidal to benzoidal structure. The experimental value of the stoichiometric coefficient is, in general, considerably lower than the theoretical one. This experience can be interpreted by means of the hot radical hypothesis.

The authors thank Zsadon Belane and Aitner Jozsefne for their participation in the research work. Orig. art. has: 2 figures and 2 tables. [JPRS: 36,862]

SUB CODE: 07 / SUBM DATE: 18Sep65 / ORIG REF: 005 / SOV REF: 001 / OTH REF: 021

Card 1/1gd

0921 0025-

L 01993-67 ENL(m)/MSP(j)/T IJP(c) WM/RM

ACC NR: AM6023688

Monograph

JR/ 33

Tudos, Ferenc

B71

An examination of the kinetics of radical polymerization based on the hypothesis of hot radicals: a survey (Rassmotreniye kinetiki radikal'noy polimerizatsii na osnove gipotezy goryachikh radikalov; obzor) Moscow, Izd-vo "Mir", 66. 0147 p. illus., biblio. Number of copies printed not given.

TOPIC TAGS: chain polymer, polymerization, radical polymerization, heat of polymerization, polymerization kinetics

PURPOSE AND COVERAGE: This book is based on the work of a Hungarian scientist, Ferenc Tudos, reported on at the Kinetic Seminar of the Institute of Chemical Physics AN SSSR. The work touches upon an extremely interesting and unresolved problem pertaining to the role played by the hot particles in kinetics. Based on the hypothesis of hot particles, it extends a large experimental material in the realm of radical polymerization. The basic premises of this work are given in the abstract presented at the meeting of the Committee on Catalysis, Hungarian Academy of Sciences, 29 May 1961. Separate parts of this communication were reported at the following conferences: Meeting on the Chemistry of Plastics (Budapest, May 1961); The Chemical Conference (Debrecen, October 1961); Conference on the Chemistry of Plastics (Smolyan, September 1962); the Chemical Conference (Seged, September 1963). This book is intended for scientists interested in the physical chemistry of polymers.

Card 1/2

UDC: 541.0/.1

L 61995-67

ACC NR: AM6023688

TABLE OF CONTENTS (abridged):

- Foreword -- 6  
Legend of basic designations -- 7  
A. General notes -- 13  
B. Stoichiometric anomalies in the case of inhibiting reactions -- 19  
C. Analysis of the measurement results and establishing of a working hypothesis -- 46  
D. Physical bases of the hypothesis of hot radicals -- 50  
E. The experimental verification of the hypothesis -- 70  
F. Generalization of the results. The role of the reaction heat in the chain reactions -- 91  
Bibliography -- 140

SUB CODE: 07, 11/ SUBM DATE: 29Dec65/ ORIG REF: 019/ SOV REF: 012/ OTH REF: 175

fv

Card 2/2

"APPROVED FOR RELEASE: 03/14/2001

CIA-RDP86-00513R001757410002-4

... by the method of polymerization and copolymerization of radical polymerization by fat-soluble

presence of the relative reactivity it was established that the elementary

APPROVED FOR RELEASE: 03/14/2001

CIA-RDP86-00513R001757410002-4"

"APPROVED FOR RELEASE: 03/14/2001

CIA-RDP86-00513R001757410002-4

APPROVED FOR RELEASE: 03/14/2001

CIA-RDP86-00513R001757410002-4"

"APPROVED FOR RELEASE: 03/14/2001

CIA-RDP86-00513R001757410002-4

APPROVED FOR RELEASE: 03/14/2001

CIA-RDP86-00513R001757410002-4"

"APPROVED FOR RELEASE: 03/14/2001

CIA-RDP86-00513R001757410002-4

APPROVED FOR RELEASE: 03/14/2001

CIA-RDP86-00513R001757410002-4"

TURCSANYI, Bela; TUDOS, Ferenc

Organic molecular compounds. Pt.2. Magy kem folyoir 71  
no.1:39-43 Ja '65.

1. Central Research Institute of Chemistry of the Hungarian  
Academy of Sciences, Budapest.

BEREZHNAYA-FEL'DESH, T.A. [Berezsnich-Foldes, T.A.]; TYUDESH, F. [Tudos, F.]

Initiated polymerization of vinyl acetate. Part 1. Vysokom.sod. 6  
no.8:1522-1528 Ag '64. (MIRA 17:10)

Initiated polymerization of vinyl acetate. Part 2. Ibid.:1529-1533

1. TSentral'nyy nauchno-issledovatel'skiy institut khimii Akademii  
nauk Vengrii, Budapest.

FOLDESNE JEREZSNICH, Tamara, kandidatus; TUDOS, Ferenc

Kinetics of radical polymerization.Pt.9. Magy kem folyoir 70  
no.5:201-207 My '64.

1. Central Research Institute of Chemistry, Hungarian Academy  
of Sciences, Budapest.

TUDOS, Ferenc

Kinetics of radical polymerization.Pt.10. Magy kem folyoir 70  
no.5:207-214 My '64.

1. Central Research Institute of Chemistry, Hungarian Academy  
of Sciences, Budapest.

TUDOS, Ferenc, a kemial tudomanyok kandidatusa

Kinetic analysis of radical polymerization of the ground of the  
hypothesis of hot radicals.Pt.2. Kem tud kozl MTA 21 no.2:191-240  
'64.

1. Central Research Institute of Chemistry, Hungarian Academy of  
Sciences, Budapest.

BICZO, Geza; LADIK, Janos; TUDOS, Ferenc; BEREZSNICH, Tamara

Calculating the atomic localization energy of polycondensed  
hydrocarbons. Magy kem folyoir 70 no. 1: 17-19 Ja '64.

1. Magyar Tudomanyos Akademia Kozponti Kemial Kutato  
Intezete, Budapest.

BICZO, G.; LADIK, J.; TUDOS, F.; BEREZSNICH, T.A.

Calculation of some atomic localization energies of various polycyclic hydrocarbons. Acta phys Hung 16 no.2:173-180 '63.

1. Central Research Institute for Chemistry of the Hungarian Academy of Sciences, Budapest.

TUDOS, F.

IRING, M. [Iring, Margit] (Budapest, XI., Fehervari) TYUDYESH, F.  
[Tudos, Ferenc], dr.; TELEKI, P. [Teleki, Piroska]

Polymerization of styrene initiated by sulphinic acid. I.  
Acta chimica Hung 35 no.3:281-296 '63.

TUDOS F.

IRING, M. [Iring, Margit] (Budapest, XI., Fehervari ut 120);  
TYUDYESH, F. [Tudos, Ferenc], dr. (Budapest, II., Pusztaszeri ut 57/69);  
TELEKSI, P. [Teleki, Piroska] (Budapest, II., Pusztaszeri ut 57/69)

Polymerization of styrene initiated by sulphinic acid. I.  
Acta chimica Hung 35 no.3:281-296 '63.

l. Chlen, Radaktsionnaya kollegiya, "Acta Chimica Academiae  
Scientiarum Hungaricae," (for Tudos).

TUDOS, Ferenc, a kemial tudomanyok kandidatusa; HEREZSNICH, Tamas; HEDER, Janos; SIMANDI, Laszlo; TURCSANII, Bela

An account of the Smolenice conference dealing with the  
chemical conversion of polymers. Kem tud kozl MTA 19  
no.2:279-282 '63.

1. Magyar Tudomanyos Akademia Kozponti Kemial Kutato  
Intezete, Budapest. 2. "A Magyar Tudomanyos Akademia  
Kemial Tudomanyok Osztalyanak Kozlemenyei" szerkeszto  
bizottsagi tagja" (for Tudos).

TUDOS, Ferenc, a kemial tudomanyok kandidatusa.

An account of my study trip to Czechoslovakia. Kem tud kozl MTA 18  
no.4:639-642 '62.

1. Magyar Tudomanyos Akademia Kosponti Kemial Kutato Intezete, Budapest,  
es "A Magyar Tudomanyos Akademia Kemial Tudomanyok Osztalyanak  
Kozlemenyesi" szerkeszto bizottsagi tagja.

HUNGARY

Budapest, Magyar Kemiai Folyoirat, Vol 69, No 8, Aug 1963, pages 371-377.

compounds. From the extent of radical concentration and the superfine structure of the EPR spectra, conclusions were drawn with regard to the electron structure of the radicals and their reactivity. The validity of the kinetic equations previously derived was also proven by means of EPR investigations. 11 Eastern European, 8 Western references.

2473  
2/2

SZPIGIN, V.I...[Spitsin, V. I.]; TUDOS, Ferenc [translator]

The present-day situation and long-range development of chemical sciences in the Soviet Union. Kem tud kozl MTA 18 no.2:173-219 '62.

1. A Szovjetunió Tudományos Akadémiaja Fizikai Kemiai Intézete, Moszkva, es Lomonoszov Egyetem, Moszkva. 2. "A Magyar Tudományos Akadémia Kemiai Tudományok Osztályának Közleményei" szerkesztő bizottsági tagja (for Tudos).

TUDOS, Ferencne

An account of my study trip to Czechoslovakia. Kem tud  
kozal MTA 19 no.2:283-285 '63.

1. Magyar Tudomanyos Akademia Kozponti Kemial Kutato  
Intezete, Budapest.

TUDOS, Tibor, okleveles villamosmernok

Automation of electric household appliances. "Villamosag" 8  
no.10;302-310 O '60.

1. Erosaramu Gyartmanyfejlesztesi Intezet.

TUDOSE, I., ing.

Construction techniques of main gas pipelines in Rumania. Petrol  
si gaze 14 no.1:28-36 Ja '63.

TUDOSE, Iulian, arh.

A good and trained mechanical servicing. Constr Buc 16 no.745:3  
18 Ap '64.

1. Seful santierului de la fabrica de incaltaminte "fregresul",  
Bucuresti al Trustului nr. 1.

TUDOSE, I.

Building and mounting activities in the petroleum industry. p. 345.

PETROL SI GAZE. (Asociatia Stiintifica a Inginerilor si Technicienilor din Romania si Ministerul Industriei Petrolului si Chimiei) Bucuresti Rumania. Vol. 10, no. 8, 1959.

Monthly list of East European Accessions (EEAI) LC Vol. 9, no.2  
Feb. 1960

Uncl.

"APPROVED FOR RELEASE: 03/14/2001

CIA-RDP86-00513R001757410002-4

TUDCSE, Mihail, ing.

Training and improving the cadres. Constr Buc 16 no.741:4  
21 Mr '64.

1. Directorul Centrului scolar de constructii-Bucuresti.

APPROVED FOR RELEASE: 03/14/2001

CIA-RDP86-00513R001757410002-4"

TUDOSE, N., ing.

Effective methods of optimum solution of directing problems  
of empty railroad cars. Rev sailor fer 11 no. 9:526-530 S '63.

1. STE.

TUDOSE, N., ing.

Operative methods for solving in the most favorable way the problem  
of directing empty railroad cars. Rev cailor fer 11 no.10:581-584  
O '63.

1. S.T.E.

"APPROVED FOR RELEASE: 03/14/2001

CIA-RDP86-00513R001757410002-4

OROS, Gh., ing.; RIZESCU, C., ing.; TUDOSE, M., ing; URсу, V., ing.

Utilization of nickel oxide in steel alloying. Metalurgia  
constr mas 15 no.8:487-490 Ag '63.

APPROVED FOR RELEASE: 03/14/2001

CIA-RDP86-00513R001757410002-4"

OROS, Gheorghe, ing.; RIZESCU, Cornel, ing.; TUDOSE, Moise, ing.;  
URSUL, Vasile, ing.

New technological methods regarding the heating of open-hearth furnaces with silica vaults. Metalurgia constr mas 15 no.6:396-402 Je '63.

1. Uzinele 23 August, Bucuresti.

FIRICA, Th., prof.; MUNTEANU, V., dr.; TUDOSE, N., dr.; ANDRONESCU, C., dr.;  
ALIONEANU, Ileana, dr.

Emergency surgery in digestive hemorrhages caused by rupture of  
esophageal or gastric varices. Med. intern. 13 no.11:1541-1548 N '61.

(HEMORRHAGE, GASTROINTESTINAL surgery)  
(ESOPHAGEAL VARICES complications)

TUDOSE, RADU

H-2

RUMANIA/Chemical Technology, Chemical Products and Their  
Application, Part 1. - Processes and Apparatus  
of Chemical Technology.

Abs Jour: Referat. Zhurnal Khimiya, No 10, 1958, 32679.

Author : Teodor Saveanu, I. Ibanescu, Radu Tudose.

Inst : Jassi Polytechnical Institute.

Title : Method of Determination of Filtration Constants at  
Filtration with Precipitate Formation.

Orig Pub: Bul. Inst. politehn. Iasi, 1956, 2, No 3-4, 129-138.

Abstract: No abstract.

Card : 1/1

"APPROVED FOR RELEASE: 03/14/2001

CIA-RDP86-00513R001757410002-4

TUDOSE, Radu Z. [Tudose, Radu S.]

Heat exchangers for fluid. Khim. i neft. mashinostr. no. 2:42-43  
(MIRA 18:4)  
F '65.

1. Yasskiy politekhnicheskiy institut, Rumynskaya Narodnaya  
Respublika.

APPROVED FOR RELEASE: 03/14/2001

CIA-RDP86-00513R001757410002-4"

"APPROVED FOR RELEASE: 03/14/2001

CIA-RDP86-00513R001757410002-4

TUDOSE, R., ing.

Column for liquid-liquid extraction. Rev. chimie Min petr 12  
no. 9:544-548 S'61

APPROVED FOR RELEASE: 03/14/2001

CIA-RDP86-00513R001757410002-4"

TUDOSE, R.  
SURNAME, Given Names

Country: Rumania

Academic Degrees: -Engineer.-

Affiliations: -not given-

Source: Bucharest, Revista de Chimie, Vol 12, No 9, Sep 1961, pp 544-548.

Data: "Column for Liquid-Liquid Extraction."

EPO 981643

"APPROVED FOR RELEASE: 03/14/2001

CIA-RDP86-00513R001757410002-4

APPROVED FOR RELEASE: 03/14/2001

CIA-RDP86-00513R001757410002-4"